



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE APPROXIMATE SUMMATION OF n TERMS OF ANY HARMONIC SERIES.

By O. L. CALLECOT, Omaha, Nebraska.

For any series $U_1 + U_2 + \dots$, we shall give notations to certain groups of successive terms; thus,

$$G_1 = U_{m+1} + U_{m+2} + \dots + U_{m+x_1}, \quad G_2 = U_{m+x_1+1} + \dots + U_{m+x_1+x_2}, \dots$$

Then for $n > m$, we have

$$(1) \quad E \equiv (G_1 + \dots + G_n) - (U_1 + \dots + U_n) = (G_1 - U_1) + \dots + (G_n - U_n),$$

$$(2) \quad E = (U_{n+1} + U_{n+2} + \dots + U_{m+x_1+\dots+x_n}) - (U_1 + U_2 + \dots + U_m),$$

since the terms U_{m+1}, \dots, U_n are common to $G_1 + \dots + G_n$ and $U_1 + \dots + U_n$.

Applying this method to the special harmonic series $1 + \frac{1}{2} + \frac{1}{3} + \dots$, and taking $m=1, x=y=w=\dots=z=3$, we have, by (1) and (2),

$$(3) \quad E = \left(\frac{1}{2} + \frac{1}{3} + \frac{1}{4} \right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} \right) + \dots + \left(\frac{1}{3n-1} + \frac{1}{3n} + \frac{1}{3n+1} \right) \\ - \left(1 + \frac{1}{2} + \dots + \frac{1}{n} \right) = \frac{2}{2 \cdot 3 \cdot 4} + \frac{2}{5 \cdot 6 \cdot 7} + \dots + \frac{2}{(3n-1)(3n)(3n+1)},$$

$$(4) \quad E = \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{3n+1} - 1.$$

$$(5) \quad \text{Hence } \frac{1}{n+1} + \dots + \frac{1}{3n+1} = 1 + \frac{2}{2 \cdot 3 \cdot 4} + \dots + \frac{2}{(3n-1)(3n)(3n+1)}.$$

But

$$(6) \quad \frac{2}{2 \cdot 3 \cdot 4} + \frac{2}{5 \cdot 6 \cdot 7} + \dots \infty = .098612 \dots$$

$$(7) \quad \frac{2}{2 \cdot 3 \cdot 4} + \dots + \frac{2}{(3n-1)(3n)(3n+1)} = .098612 \dots - \frac{1}{3(3n+1)(3n+2)} + \delta,$$

$$\delta < \frac{2}{(3n+1)(3n+2)(3n+3)(3n+4)}.$$

Now any number which gives the remainder 1 on division by 3 may be taken as the last in a group of $2n+1$ terms. We may therefore arrange the series $1 + \frac{1}{2} + \dots$ in successive groups of $2n+1$ terms by throwing out 0, 1, or 2 terms for each group, as the case may be.

$$(8) \quad \text{Hence } 1 + \frac{1}{2} + \dots + \frac{1}{n'} = G + D + U,$$

where G is the number of groups in n' terms, U is the sum of the deleted terms, and D is the sum of the difference groups, $\frac{2}{2.3.4} + \dots + \frac{2}{(3n-1)(3n)(3n+1)}$. If n is not taken smaller than 15,

$$(9) \quad D = (.098612\dots\dots)G - \frac{10^G - 1}{27(10^{G-1})(3g+1)(3g+2)} - \delta,$$

$$\delta < \frac{1}{216(3g+1)(3g+2)},$$

where g is the n of the group of lowest denominator.

Since the formula applies to the series $\frac{1}{r} + \frac{1}{r+1} + \dots$, $r > 15$, we apply the following table for the sum of the earlier terms.

TABLE GIVING SUM OF FIRST n TERMS OF SERIES $1 + \frac{1}{2} + \dots$

n	Sum	n	Sum
2	1.5	23	3.734291
3	1.833333	24	3.775958
4	2.083333	25	3.815958
5	2.283333	26	3.854419
6	2.449999	27	3.891456
7	2.592857	28	3.927171
8	2.717857	29	3.961653
9	2.828968	30	3.994987
10	2.928968	31	4.027245
11	3.019877	32	4.058495
12	3.103210	33	4.088798
13	3.180133	34	4.110209
14	3.251562	35	4.138781
15	3.318228	36	4.166559
16	3.380728	37	4.193586
17	3.439552	38	4.219901
18	3.495108	39	4.245542
19	3.547739	40	4.270542
20	3.597739	41	4.294933
21	3.645338	42	4.318742
22	3.690813	43	4.341998

Example. Find the sum approximately of $1 + \frac{1}{2} + \dots + \frac{1}{10,000}$.

	10000	<i>Denominators of terms in U.</i>	
3)	9999	0	0
	3333	3333	3332
	<hr/>		
	3331		
3)	3330		
	1110	1110	1109
	<hr/>		
	1108		
3)	1107		
	369	369	368
	<hr/>		
	367		
3)	366		
	122	122	0
	<hr/>		
	121		
3)	120; ———40		

$$B=4.270542$$

$$U=0.016088$$

$$(1.098612)5=5.493061$$

$$\underline{9.779692}$$

$$\text{Second term of (9)} = -0.000025$$

$$\text{Sum} = 9.77966 \dots - \delta, \delta < .000,000,313.$$

Any harmonic series $\frac{1}{a} + \frac{1}{a+d} + \frac{1}{a+2d} + \dots$ may be reduced to the form

$$\frac{1}{d} \left(\frac{1}{n+f} + \frac{1}{n+1+f} + \frac{1}{n+2+f} + \dots \right)$$

where f is the fractional part of the result of dividing a by d , and n is the integral part of the result. Also,

$$\begin{aligned} & \frac{1}{d} \left(\frac{1}{n} + \frac{1}{n+1} + \dots + \frac{1}{n'} \right) - \frac{1}{d} \left(\frac{1}{n+f} + \frac{1}{n+1+f} + \dots + \frac{1}{n'+f} \right) \\ &= \frac{1}{d} \left(\frac{f}{n(n+f)} + \frac{f}{(n+1)(n+1+f)} + \dots + \frac{f}{n'(n'+f)} \right). \end{aligned}$$

Since the value of $\frac{1}{d} \left(\frac{1}{n} + \dots + \frac{1}{n'} \right)$ may be found according to methods already

given, and since the series $\frac{1}{d} \left(\frac{f}{n(n+f)} + \dots \right)$ converges rapidly, we may approximately determine the sum of n terms of any harmonic series.

NOTE ON THE MAXIMUM INDICATOR OF CERTAIN ODD NUMBERS.

By REV. R. D. CARMICHAEL, Hartselle, Alabama.

If p , the least prime factor of N , is of the form $4l+1$, the maximum indicator of N is a multiple of 4.

In the MONTHLY, May, 1905, p. 107, I have shown that, for p a prime of the form $4l+1$, we have

$$\left(1.2.3 \dots \frac{p-1}{2} \right)^2 \equiv -1 \pmod{p}.$$

Hence, if $\left(1.2.3 \dots \frac{p-1}{2} \right)^{2n} \equiv 1 \pmod{N}$, n is even; and therefore the maximum indicator of N , being a multiple of the least value $2n$ satisfying the above congruence, is also a multiple of 4.

Corollary. *The equation $y^4 = mx + 1$ has at least one positive integral solution when the least factor of m is congruent to unity modulo 4.*

By Wilson's theorem, it is easily shown as above that

The maximum indicator of any odd number is even.

Corollary. *The equation $y^2 = mx + 1$ has at least one positive integral solution, when m is odd.*

If p and $2p-1$ are odd primes, the maximum indicator of $p(2p-1)$ is a multiple of 4.

As in the first congruence above we have

$$(1.2.3 \dots \overline{p-1})^2 \equiv -1 \pmod{2p-1}.$$

By Wilson's theorem,

$$(1.2.3 \dots \overline{p-1})^2 \equiv 1 \pmod{p}.$$

$$\text{Thus } (1.2.3 \dots \overline{p-1})^4 \equiv 1 \pmod{p \cdot 2p-1}.$$

Now, $(1.2.3 \dots \overline{p-1})$ is not congruent to 1 modulo $p \cdot 2p-1$. It is thus shown that one indicator, at least, is 4. Hence the maximum indicator is a multiple of 4.